

日期

2019年09月22日

內容摘要：

(填寫說明：

1. 如有附件請註明，如簡報檔、全文檔等
2. 需有問題與討論：請註明姓名並包含醫學倫理及 EBM 之應用
3. 需有總結，請註明做結論者【主持人】姓名
4. 請自行編排頁碼)

六大核心能力

- 病人照護
- 人際溝通技能
- 醫學知識
- 從工作學習及成長
- 專業素養
- 制度下之臨床工作

B2 月新團會議室

9/22 0830 - 0930

Case conference

Present: R. 蔡宇承

(Q/A) Supervisor: F2 徐英洲

(EBM)

Q (F2 徐英洲): Canadian C7 head rule Dangerous mechanism

A (R. 蔡宇承): Amnesia before impact, 20 min. GCS < 15 at 2 hrs after injury. Suspect open or depressed skull base fx. Any sign of basilar skull fx, vomiting > 2 episode. Age > 65y

Q (F2 徐英洲): Head C7 rule (Canadian). Dangerous mechanism

A (R. 李怡穎): pedestrian, occupant ejected, fall from elevation (23 feet on 5 stairs)

Q (F2 徐英洲): Canadian head C7 rule 不適用

A (R. 蔡宇承): Nontrauma cases. GCS < 13. Age < 16 year. Coumadin or bleedn. disorder. obvious open skull fx

Q (F2 徐英洲): Sign of basilar skull fx

A (R2 施廣泰): Hemotympanum, Raccoon eyes. CSF otorrhea/Rhinorrhea Battle's sign

Q (F2 徐英洲): Ottawa ankle rule

A (R. 林哲羣): Bone tender over distal 6cm of posterior edge of tibia or top of medial malleolus

o An inability to bear weight both immediately and in the emergency department for four step

Q (F2 徐英洲): Ottawa foot rule

A (R2 吳冠蓉): Any pain in the midfoot zone Bone tender at the base of 1st metatarsal or navicular bone

不能 weight bearing for 4 step

內容摘要 (續):

Q(王維英洲): lower triad fx

A(R.劉邦民): Calcaneal comminuted fx.

Brust fx of lumbar or thoracic fx

Q(F.維英洲): 屈股著地帶小以那裡骨折

A(R.鄭凱文): Thoracolumbar fx

Q(F.維英洲): Monteggia fx

A(R.蔡可威): Fx of proximal third of ulna

+ Dislocation of head of the radius

Q(F.維英洲): Galeazzi's fx

A(R.蔡守承): Fracture of radius with dislocation of distal ~~radius~~ radioulnar joint

VS Comment: Trauma 病人 比 要 詳 盡. from Head to toe.
F. 維英洲 該 切 的 時 還 是 要 切 (Canadian Head of rule)

紀錄者: 吳邦民

